METEOROLOGICAL OBSERVATIONS


STATION NOTES

The climatological readings for Kingston are made from several stations. Temperatures and humidity are measured in a Stevenson Screen situated at St. George's College and supervised by the Society of Jesuits. The exposure is good and the readings are considered representative of the area. Rainfall is measured at the Public Works Office but there are several other gauges within a two-mile radius with which to compare. This is necessary with the capricious nature of Jamaica rain. Cloud weather and wind are recorded at Palisadoes Airport situated on the seaward side of Kingston Bay. This is a full time synoptic and climatological meteorological station, first class, maintained by Air Ministry staff, and is the forecast centre for the western Caribbean. Here the exposure is excellent but the cloud amount tends to be less than over the City. Showers are also less frequent. The day sea breeze sets in earlier at the Airport and apparently is stronger, while the night land breeze which cools the urban area does not appreciably affect the Airport till dawn. Wind is recorded by Dines Pressure Tube Recording Anemometer.

Both Morant Point and Negril Point Lighthouse are full synoptic reporting stations with a good complement of instruments including Cup Anemometers. Exposures are open and very good, the sites at the extreme east and west extremities of the Island being excellent for their main purpose of weather reporting for immediate international use.



QC 987 ,J25 M48 1949



National Oceanic and Atmospheric Administration

Climate Database Modernization Program

ERRATA NOTICE

One or more conditions of the original document may affect the quality of the image, such as:

Discolored pages
Faded or light ink
Binding intrudes into the text

This document has been imaged through the NOAA Climate Database Modernization Program. To view the original document, please contact the NOAA Central Library in Silver Spring, MD at (301) 713-2607 x124 or www.reference@nodc.noaa.gov.

LASON Imaging Subcontractor 12200 Kiln Court Beltsville, MD 20704-1387 March 28, 2002

KINGSTON, JAMAICA.--Mean Meteorological Results for the Year 1949. Longitude 76°48' West Latitude 17°58' North

			_	₹;	Air Temperatures Fah.	eratur h.	8 9				Dew	Dew Point		Rela	Relative Humidity		Amount of Cloud Tenths	oud the		Rainfall Inches			No.	Weather No. of days of	s of			ŝ	. of 0	Wind	Wind No. of Observations	of	
	PARILL			•	Means of		Absolu	te Ma Min.	Absolute Max. and Min.					l							1		torm	1	t	'		<u> </u>		1			<u> </u>
Month	Mean Pre	н. Т.н.	— Н Меви		Min. M	Мах. Л	Min.	Date	Max.	Date es	1- 13. II.	es Ej	nashi a	1- E	⇔ ⊟ Mean	ei	7 m.p.m.	n.m.	LatoT	Max.	Date	Rain	З тэрпийТ	Clear Sky	dasonevO	aslaD	<u>.</u>	ਬ <u>ਂ</u> ਤ	80 ज	vi	≥	z ≽	S S anlaO
Jan. 30.	30.007 69	.2 84.6	.6 75.9		67.0 84.	90	64.6	19 9	91.4	- e	 8	70 67			61 77	<u>01</u>	9 3.1	23 23	0.30	0.22	8	က	C	12	0	0	16	2 12	6	122	2	9	1 1
Feb. 30.	30.025 69	69.7 85	85.4 77	79 67	7.4 87	ī.	65.0	17 9	90.3	17 6		02		91	59 79	્ય	9.2.9	52	0.02	0.02				83		- 0	 	3 19	19	7	_	9	9 11
Mar. 29.	29.956 71	71.1	83.8	نء	69.1 87		65.0	11 9	91.0	18		69 67		9 08	61 73	<u>€4</u>	0 3.0	62	0.25	0.19	- 22	4	0	27	0		 	1 12	83	17	_	ಌ	5 27
Apr. 29.	29.969 76	76.7	- 4		72.0 88	<u>. 6,i</u>	67.3		93.7	- 92	2 29	11 69		73 6	61 71	2.7	7 3.3	2.9	0.29	0.22	- 63	~~		24	0	0	=======================================	4 27	- E3	-	:		6 14
May 29.	29.930 76	76.9	.3 80.	0	72.9 87.	r-	* 0.07	*17	89.2	13	<u></u>	74 72	r	79 7	70 7	<u>~</u>	6 4.7	7.	2.61	0.93	19	~	9	6	c 4		16 4	4 13	15	:	:		19 20
June 29	29.959 77	77.9 85	85.0 81	0.	74.3	88.8	71.3	14	93.5	-22	2	73 71			68 76		4.5	4.7	99:0	0.33	~	<u>.</u>	6	2				<u> </u>	8	;	:		10 13
July 29	29.973	77.7	85.1		74.5 90	90.1	74.7	01	92.9	200		74 73		78	70 78	<u></u>	7 4.0		2.82	2.13		9	133	12	0	0				က	:		19 16
Aug. 29	29.949 77	98 82	85.9 81	81.6	74.5 9(90.2	72.2	08 08	94.1	4		75 73			69 77	<u> </u>	7 3.9	4.0	1.37	0.53	8	9	=	10	0	0		8	37	4	-	63	13 20
Sept. 29	29.904 77	77.3 85	85.4 81	81.3	74.9	89.4	72.2	-54	93.0	- × - × - ×		57 27		82 _ 7	71 81	4	5.4.3	5.0	9.15	3.70	- 23	=	17	∞	_	0		2 26	**	ಌ	က	63	13 27
Oct. 29.	29.893 75	75.9 85.	2	80.4	73.9	88.7	9.02	24	93.7	29 7		74 73		85 7	71 81	<u> </u>	7 3.6	. .	2.23	0.71	=======================================		15	.: ::	0		14	2 15		9	60	*1117	19 32
Nov. 29	29 903 74	74.4 84	84.5 79	79.3 72.	2.6 87	70	0.99	5 62	90.0	11 /		73 71		87 6	88 81	41	5 4.7	600	2.13	1.10	- 18	∞	9	01		0 1	17 2	18	14	77	C3	:	18 35
Dec. 29.	29.923 70	8 81	.2 76.	.5 69	9.5 86	90	65.3	16 9	90.5	10	2 29	72 69		72	£7 88	<u>. 64</u>	-7- -3- -3-	1.9	0.75	0.24	8	9	m	61	_	0	 %	 	6	51	9	727	24 35
Means 29.	.950 74	74.6 84.	7 79		71.9	88.1	85 65	- Gs :	91.9	9	69	17 87		.9	67 78	2. E.	80	3.5	:	:	:	:	:	:	:	<u> </u>	 		:	:	<u>† :</u>	:	1 :
Totals							:	-	:		 :	 :	 -				<u>:</u>		22.58	:	:	67	8	196	9	0 203		270	348	æ	3	20	180/060

*And any other days.

*Imporation to the Standard of National Physical Laboratory 32°, Gravity at Latitude 45°, and to Mean Sea Level. The 75th Meridian West of Greenwich Standard Time used. Altitude of Standard Barometer above Mean Sea Level 24 ft. Height of Thermometers alove ground 4 ft. of Rain-guage above ground 3 ft. Height of Anemometer above ground 50 ft.

*And on other days.

Glear Sky--0-2/8 Cloud.

Overcast. -28 Cloud.

Overcast. -28 Cloud.

Overcast. -28 Cloud.

However to the Standard of National Physical Laboratory 32°. Gravity at Latitude 45° and to Mean Sea Level. The 75th Meridian West of Greenwich Standard Time used. Attitude of Standard Barometer above for Standard of Thermometers above ground 4 ft. 6 inches. Height of Rain-gauge above ground 6 ft. 5 ins. Height of Anemometer above the ground 9 ft. to the 31st May, 1936, and 22 ft. above ground from 1st June, 1936.

MORANT POINT LIGHTHOUSE, JAMAICA.—Mean Meteorological Results for the Year 1949. Longitude 76° 12' West Latitude 17° 56' North

		1	-	Air	Air Temperatures Fah.	tures	ļ			Ã	Dew Point		Hun	Relative Humidity	<u> </u>	를 하는 -	Amount of Cloud Tenths		Rainfall Inches	_		Weather No. of days	Weather of days	jo			S. No.	of Obs	Wind Observations	B of		
911188				Ž	Means of		solute M	Absolute Max. and Min.	and							named Photograph						mios	_			į	E	5 5	1	ŧ		
Mean Pre	- E	p.m.	Мевп	Min.	l. Max.	. Min.	: Date	Мах.	Date	1- 1. H.	P. II.	7.0.0	a.m.	P.m. 7	7.m.c	r- H	1 7 mr. 9.m	letoT	Max.	Date	niaH	ThundT	Clear Sky	Overcast Gales		á c					호 조	Calm
30.021	21 76.5	.5 79.9	9 77.8	E	8 81.1	- 69	-2	<u>æ</u>	31	66	72	71	-19	- &	9 18	5 6.	1 3.8	3.53	1.58		17	:		12	19	88	32	62	0	67	63	-
30.054	54 76.1	.1 81.0	-26	9 73.	88 88	-2 -83	- 81	88	21	66	11	22	28	 98	₹ 5	_ -6	- 63 - 63 - 63	1.42	0.38		on on	:	:			88	83	 	0 0		-	0
8	.961 74.5	.5 82.0	0 77.4	4 71.3	3 84.1			\$	15	20	72	8	33	71	79	<u> </u>	0 2.5	1.77	7 0.70	- 88	2	:	:	- . :	12	8	27	19	0 0	က	6.7	0
29.6	.976 78.	0 83.6	8 78.7	7 75.0	0 85.1	1 67		%	ž.	23	7.5	23	88	- 47	88 4	*- 	9.2.4	0.74	0.35		6	:			4	13	6		0 0	0	-	
29.932	28	- 4-	8.3	3 75.	2 84.3		- ₹	- 8	7	74	92	7.	æ	8 62	86 4.	4	2 4.1	16.20	4.32	- 10	82	rO.			~	63	12		- 0		0	ო
85 83	.964	88	8 80.1	29	25	17 6.		88	73	35	76	22	25	82	- 2 -	∞	.5 3.5	6.47	-3.23		==	4		13	-	•	19		2 0	0	0	
86	980 81.0	0 84.4	81.0	0 78.1	1 86.4		17	8	19	76	1	76	28	<u>8</u>	- 7.		1 3.2	38.	1.38	8	13	9				90	क्ष		2 0	٥		0
28.950	83	3 84.8	8 81.6		8.9	02	11		ន	75	F	-22	2		84 3.1	<u> </u>	6 3.5	11.23	3 2.62	8	16	2				۲-	8 2		0 1		0	C4
29.897	88	3 84.7	9.18	*	.5 86.7	7 7	9.	8	ţ	92	28	7.8	\$6	- 8°	88 83	86 23	12) 60	18 13	85 85	92	8	91		11 2	188	9	21	45	0 2		-	
88	892 79.5	5 84.7	81.1	7.	2 85.9	67		87	7"	75	78	22	62	- 62	82	86 89	9.2	10.91	2.25	=	22	٢			15	z	6			**		70
29.883	13 78.9	9 83.0	79.8	74.6	3 24.5	8	9	88	7	74	92	75	138	-8-	26 4.		9.8	80.6	1 2.63		19	64		: :	*	R	r ~	23	1 0	4	7-4	
29.927	78.0	80.8	78.6	74.5	88.25	3	22	87	36	22	73	72	28			<u></u>		3.10	1.14	 	135		:	en-	37	\$	13	~~			0	0
Means 29.954	4 78.6	0.88	79.4	74.9	7.78	*	:	126	:	23	92	174	88	8 82	8	3 4.1	1 3.2	86.49	:	ļ :		:	:		<u> :</u>	:	1 :	-	:	:		:
:	:	;		:																		-										

*And on other days.

†Clear Sky — 0-2/8 Cloud.

10vercast —8/8 Cloud.

10vercast —8/8 Cloud.

10vercast —8/8 Cloud.

Barometer Pressure reduced to the Standard of National Physical Laboratory 32°. Gravity at Latitude 45° and to Mean Sea Level. The 75th Meridian West of Greenwich Standard Time used. Altitude of Standard Barometer above Mean Sea Level 8 ft. Height of Thermometers above ground at 4 ft. 6 ins. Height of Rain-gauge above ground 3 ft. Height of Anemometer above ground 18 ft.

THE WEATHER OF 1949

The rainfall for the Island was 88% of the 70-year Mean. The parishes of St. Thomas (116%) and Hanover (105%) received more rain than normal, whilst the Parishes of St. Mary (74%), Portland (73%), Clarendon (74%) and Kingston (70%) were well below average. The rainfall for the remaining Parishes was about normal. On examining the island mean figures month by month, May was the wettest month in 1949; the Parish of St. Thomas had 227% of the normal, whilst St. Ann, St. Elizabeth and Westmoreland received almost half as much again as normal. September and December were rather wetter than usual and November in some areas was above average, particularly in St. Thomas (188%) and Clarendon (151%) in September, and Hanover (163%), St. James (155%) and Westmoreland (174%) in November, and St. Andrew (216%) in December. January and February were very dry in most Parishes. Kingston only received 17% in January and 3% in February of its normal rainfall.

Temperatures were a little above average for the year but nights were cooler than normal in January, February, March and December.

Extremes—During January, 23 stations reported no measurable rain, whilst 22 stations also received no rainfall.

The highest recorded temperature in Kingston was 94.1°F. on 4th August—the lowest Maximum since 1938. The coolest night occurred on 19th January when the mercury fell to 64.6°F. At Hope, the highest temperature was 96.4°F. in September, whilst January, February and March were the coolest months for at least ten years.

Northers—Light Northers penetrated as far as Jamaica in January, May and December, but there were no reports of high winds to damage crops. The weather situation in May was similar to that of May 1948 but fortunately, the disastrous floods of that year did not again occur.

Tropical Storms—There were eleven (11) tropical disturbances in and around the Caribbean in 1949, but fortunately, none came near enough to Jamaica to warrant the issue of a warning for the Island.

It was not until August 21st, that the first storm of the year was discovered, some 300 miles North of Puerto Rico. It moved WNW and then slowly curved North to pass over a Lightship off Cape Hatteras. The first storm in the Caribbean proper was located near Martinique on August 30th, but by September 2nd, the storm had weakened so much that no definite centre could be found. The sixth storm of the season and the second in the Caribbean was located 100 miles SSE of St. Croix on September 21st. It moved WNW to reach the south-east coast of the Dominican Republic and then fortunately dissipated, though not before 15 lives were lost there. The last hurricane of the season was located by aircraft 50 miles East of Swan Island on November 3rd, but fortunately, for Jamaica, drifted slowly SSW until it reached Honduras, where it dissipated rapidly.